

Leading Causes of Death*, by Sex,** Maine, 2009

Cause of Death		Number of Deaths	Crude Rate		Age-adjusted Rate	
			Rate	95% CI	Rate	95% CI
Males						
1	Malignant neoplasms (C00-C97)	1661	260.9	(248.4-273.5)	235.6	(224-247.1)
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1375	216.0	(204.6-227.4)	207.2	(196.1-218.4)
3	Chronic lower respiratory diseases (J40-J47)	387	60.8	(54.7-66.9)	58.3	(52.4-64.1)
4	Unintentional injuries (V01-X59,Y85-Y86)	379	59.5	(53.5-65.5)	58.2	(52.2-64.2)
5	Cerebrovascular diseases (I60-I69)	244	38.3	(33.5-43.1)	38.1	(33.3-42.9)
6	Diabetes mellitus (E10-E14)	188	29.5	(25.3-33.8)	26.2	(22.4-30)
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	155	24.3	(20.5-28.2)	23.2	(19.5-27)
8	Alzheimer's disease (G30)	147	23.1	(19.4-26.8)	24.7	(20.7-28.7)
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	128	20.1	(16.6-23.6)	19.9	(16.4-23.4)
10	Influenza and pneumonia (J09-J18)	106	16.7	(13.5-19.8)	16.4	(13.3-19.6)
11	Chronic liver disease and cirrhosis (K70,K73-K74)	79	12.4	(9.8-15.5)	10.5	(8.3-13.2)
12	Parkinson's disease (G20-G21)	71	11.2	(8.7-14.1)	11.6	(9-14.6)
13	Pneumonitis due to solids and liquids (J69)	60	9.4	(7.2-12.1)	9.8	(7.5-12.7)
14	Septicemia (A40-A41)	56	8.8	(6.6-11.4)	8.1	(6.1-10.6)
15	Aortic aneurysm and dissection (I71)	46	7.2	(5.3-9.6)	7.2	(5.2-9.6)
16	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	6.4	(4.6-8.7)	6.1	(4.4-8.3)
17	Essential hypertension and hypertensive renal disease (I10,I12,I15)	40	6.3	(4.5-8.6)	5.7	(4-7.8)
18	Enterocolitis due to Clostridium difficile (A04.7)	24	3.8	(2.4-5.6)	3.8	(2.5-5.7)
19	Viral hepatitis (B15-B19)	23	3.6	(2.3-5.4)	2.8	(1.7-4.2)
20	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	DNR	(1.8-4.7)	DNR	(1.6-4.4)

ICD-10 codes are grouped by: ICD-10 113 Cause List

Crude rates are deaths per 100,000 population

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population

95% CI: 95% confidence interval of the rate.

Data Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death 1999-2009 on CDC WONDER Online Database, released 2012.

DNR: Data not Reliable. Death rates based upon a numerator of 20 or less are suppressed by the National Center for Health Statistics and are reported in this table as "Unreliable."

Leading Causes of Death*, by Sex,** Maine, 2009

Cause of Death		Number of Deaths	Crude Rate		Age-adjusted Rate	
			Rate	95% CI	Rate	95% CI
Females						
1	Malignant neoplasms (C00-C97)	1472	220.4	(209.1-231.6)	158.6	(150.4-166.9)
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1296	194.0	(183.5-204.6)	124.2	(117.3-131.1)
3	Chronic lower respiratory diseases (J40-J47)	434	65.0	(58.9-71.1)	44.8	(40.6-49.1)
4	Cerebrovascular diseases (I60-I69)	403	60.3	(54.4-66.2)	38.6	(34.7-42.4)
5	Alzheimer's disease (G30)	319	47.8	(42.5-53)	29.0	(25.8-32.2)
6	Unintentional injuries (V01-X59,Y85-Y86)	218	32.6	(28.3-37)	25.5	(21.9-29)
7	Diabetes mellitus (E10-E14)	162	24.3	(20.5-28)	17.0	(14.4-19.7)
8	Influenza and pneumonia (J09-J18)	144	21.6	(18-25.1)	13.4	(11.2-15.7)
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	144	21.6	(18-25.1)	14.3	(11.9-16.7)
10	Septicemia (A40-A41)	76	11.4	(9-14.2)	7.9	(6.2-10)
11	Chronic liver disease and cirrhosis (K70,K73-K74)	65	9.7	(7.5-12.4)	7.6	(5.8-9.7)
12	Parkinson's disease (G20-G21)	56	8.4	(6.3-10.9)	5.5	(4.2-7.2)
13	Pneumonitis due to solids and liquids (J69)	45	6.7	(4.9-9)	4.3	(3.1-5.7)
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	43	6.4	(4.7-8.7)	4.1	(2.9-5.5)
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	6.3	(4.5-8.5)	5.9	(4.2-8.1)
16	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	5.4	(3.8-7.5)	3.7	(2.6-5.1)
17	Enterocolitis due to Clostridium difficile (A04.7)	35	5.2	(3.6-7.3)	3.2	(2.2-4.5)
18	Aortic aneurysm and dissection (I71)	33	4.9	(3.4-6.9)	3.4	(2.3-4.7)
19	Anemias (D50-D64)	13	DNR	(1-3.3)	DNR	(0.6-2.2)
20	Atherosclerosis (I70)	11	DNR	(0.8-2.9)	DNR	(0.5-1.8)

ICD-10 codes are grouped by: ICD-10 113 Cause List

Crude rates are deaths per 100,000 population

Age-adjusted rates are deaths per 100,000 population age-adjusted to the U.S. 2000 standard population

95% CI: 95% confidence interval of the rate.

Data Source: Centers for Disease Control and Prevention, National Center for Health Statistics. Underlying Cause of Death 1999-2009 on CDC WONDER Online Database, released 2012.

DNR: Data not Reliable. Death rates based upon a numerator of 20 or less are suppressed by the National Center for Health Statistics and are reported in this table as "Unreliable."